

# Remeflox C Vet Bolus

Ciprofloxacin Hydrochloride BP

**Composition:** Each Bolus contains Ciprofloxacin Hydrochloride BP equivalent to Ciprofloxacin 750 mg.

**Description :** Remeflox C (Ciprofloxacin) is a synthetic fluoro-quinolone derivatives antimicrobial agent having broad spectrum activity.

**Mode of action :** Remeflox C (Ciprofloxacin) is a 2<sup>nd</sup> generation fluoroquinolone and broad-spectrum bactericidal antibiotic. Ciprofloxacin kills the bacteria by inhibiting DNA gyrase (Topoisomerase-II) enzyme.

## **Indication:**

**Livestock:** Remeflox C (Ciprofloxacin) Bolus is indicated for the treatment of diseases such as Respiratory tract infection, Urinary tract infection, Gastrointestinal tract infection, Skin & soft tissue infection caused by both gram positive and gram negative bacteria (*Actinobacillus*, *Pasteurella*, *Streptococcus*, *E.coli*, *Bordetella*, *Salmonella* and also *Mycoplasma*).

**Poultry:** Remeflox C (Ciprofloxacin) Bolus is indicated for the treatment of disease caused by both gram positive and gram negative bacteria, like *Colibacillosis*, *Salmonellosis*, *Infectious Coryza*, *Mycoplasmosis (C.R.D)*, *Streptococcosis*, *Staphylococcosis*, *Pasteurellosis*, Secondary bacterial infection in case of viral diseases, such as Gumboro, Ranikhet etc.

## **Dosage & Administration:**

**Poultry:** Dissolve 1 Bolus 5-7 Liter of water & administer daily for 3-5 days.

**Cattle, Sheep, Goat & Other animals:** Administer 1 Bolus /50 Kg body weight daily for 3-5 days.

Or, as prescribed by the registered veterinary doctor

**Contra-Indication:** Ciprofloxacin should not be used together with the Chloramphenicol, Macrolides or Tetracycline.

## **Withdrawal time:**

**Meat:** 12 days, **Milk and Egg:** 03 days.

**Storage:** Protect from light, store in a cool & dry place and keep out of reach of children.

**Presentation:** Each box contains 5 X 6 Bolus in Blister Strips.

