

Remetrim Vet Bolus

Sulphadiazine BP & Trimethoprim BP

Composition: Each Bolus contains Sulphadiazine BP 1,000 mg & Trimethoprim BP 200 mg

Mode of action: Remetrim Bolus acts as competitive inhibitor in the first step of para-aminobenzoic acid (PABA) synthesis of bacteria. Trimethoprim inhibits the second stage of PABA synthesis. Sulphadiazine and trimethoprim show synergistic action of inhibition of PABA synthesis. As a result, the efficacy of Remetrim Bolus increases several times.

Indication: Remetrim Bolus is used for the prevention and treatment of different infections caused by different gram- positive & gram- negative bacteria in livestock. It is also used in dermatitis and wound infections. Remetrim Bolus is also used for the treatment of Hemorrhagic septicemia and Diphtheria of calf, Pasteurellosis, Actinomycosis, Actinobacillosis, Coccidiosis, Mastitis, Metritis, Necrotic Rhinitis, Keratitis infection, Polyarthrititis, oral necrobacillosis, Foot rot, Pneumonia, Salmonellosis, Toxoplasmosis, Bronchitis, Bronchopneumonia etc.

Dosage & Administration:

For oral administration: Smashed Remetrim Bolus can be administered via feed or drinking water

Cattle, Horse, Sheep & Goat: 1 Bolus per 40 kg body weight daily or 30 mg per kg body weight daily (use Remetrim Bolus twice a day by giving half of the dose).

Intrauterine: Smashed Remetrim Bolus can be used via mild hot water by a catheter.

Cow, Buffalo and Horse: 2-4 Bolus

Sheep & Goat: 1-2 Bolus.

Or, as prescribed by the registered veterinary doctor.

Contra-Indication: Remetrim Bolus is contraindicated in animal with known hypersensitivity to active ingredients.

Precaution: In case of hypersensitivity, adequate fluid intake should be ensured during therapy.

Side effect: At the recommended dose, side effects are rare and mild. Sometimes for prolonged uses hematological reactions, agranulocytosis and Vitamin K deficiency may occur. In some cases crystalluria may occur.

Withdrawal Time:

For meat: Cattle, Sheep & Goat : 8 days.

Storage: Protect from light, store in a cool & dry place and keep out of reach of children.

Presentation: Each box contains 4 X 6 Bolus in Blister Strips.