

# Bentronil® vet Injection

Nitroxinil BP

**Composition:** Each ml contains Nitroxinil BP 340 mg.

**Indications:** Bentronil Injection is indicated for the effective treatment and control of the infestations of mature and immature liver flukes (*Fasciolagigantica*, *Fasciola hepatica*), harmful species of gastrointestinal roundworms such as *Haemonchuscontortus*, *Bunostomum*, *Oesophagostomum*, *Oestrusovisand parafilaria*, *Syngamus* of birds and *Ancylostoma* of dogs.

## Dosage & Administration:

Bentronil Injection should be injected only by subcutaneous route.

**Cow, Buffalo:** 1.5 ml per 50 kg of body weight.

**Sheep, Goat:**

Body Weight	Dose	Body Weight	Dose
14-20 Kg	0.5 ml	41-55 Kg	1.5 ml
21-30 Kg	0.75 ml	56-75 Kg	2.0 ml
31-40 Kg	1.0 ml	>75 Kg	2.5 ml

Or, as prescribed by the registered veterinary doctor.

**Contraindication:** It should not be used in animals with a history of hypersensitivity to the active ingredients. Do not use in animals producing milk for human consumption.

**Side Effects:** No side effects have been observed at recommended doses.

**Drug Interaction:** In compatibility studies with other drugs, Bentronil Injection was found to be safe when administered concurrently with antinematodal drugs (e.g. Thiabendazole, Phenothiazine, Naphthaophos, Tetramisole etc.)

**Precaution:** Bentronil Injection is not for intravenous or intramuscular use.

## Withdrawal period:

Meat: 30 days.

Milk: 5 days.

**Storage:** Protect from light, store in a cool & dry place and keep out of reach of children.

**Presentation:** 10 ml, 25 ml, 30 ml, 50 ml and 100 ml Vial.